INDEX TO VOLUME 79

New scientific names are printed in bold face

- Abrahamson, Warren G., and Kenneth D. McCrea, Ultraviolet light reflection and absorption patterns in populations of Rudbeckia (Compositae) 269-277
- Acer saccharum 123
- Adenocalymma inundatum 438
- Allenrolfea mexicana Lundell (Chenopodiaceae): Its conspecificity with A. occidentalis (S. Wats.) Kuntze 130-132
- American Bignoniaceae, Notes on middle 430-444
- Amphitecna 430-444; apiculata sp. nov. 435. breedlovei sp. nov. 432. costata sp. nov. 431, latifolia 438, montana 437, silvicola 437, steyermarkii 436
- Aquatic plants in New England, Observations on some uncommon 445-451
- Arkansas, Morphological intergradation of varieties of Bidens aristosa (Compositae) in Northern 203-213
- Arrabidaea costaricensis 439, podopogon comb. nov. 439, pubescens 440, verrucosa 440
- Bahamas reexamined, Vernonia (Compositae) in the 147-159
- Baranov, A. I., Preliminary notes on the structure of stigmatic surfaces in the Begoniaceae 283-287
- Baranov, A. I., Rudbeckia serotina f. tubuliformis in New England 165
- Bassett, I. J., See McNeill, J. and C. W. Crompton
- Battersia mirabilis, A crustose sphacelarioid new to northeastern North America 292-299
- Begoniaceae, Preliminary notes on the stigmatic surfaces in the 283-287
- Bernhardt, Peter, The pollination ecol-

- ogy of a population of Erythronium americanum Ker. (Liliaceae) 278-282
- Bidens aristosa (Compositae) in Northern Arkansas, Morphological intergradation of varieties of 203-213
- Bidens aristosa 203-213; differences between varieties of 204; various characteristics of population 1 207 & 210; various characteristics of population II 208 & 211
- Bignoniaceae, Notes on Middle American 430 444
- Biosystematics of Psilostrophe DC. (Compositae) I. Chromosome variability 169-189
- Brown, Roy Curtiss, Biosystematics of Psilostrophe DC. (Compositae) I. Chromosome variability 169-189
- Burk, C. John and Scott D. Lauermann, Catalpa speciosa naturalized in western Massachusetts 305-307
- Canne, Judith M., A revision of the genus Galinsoga (Compositae : Heliantheae) 319-389
- Catalpa speciosa naturalized in western Massachusetts 305-307
- Ceratophyllum tetragonolobum 440 Champlin, Richard L., Red form of Lindera benzoin 166
- Chromosome studies in North and Central American species of Pectis L. (Compositae : Tageteae) 79-94
- Chromosome studies of Parnassia and Lepuropetalon (Saxifragaceae) from the eastern United States. A new base number for Parnassia 573-578
- Chromosome variability. Biosystematics of Psilostrophe DC. (Compositae)
 I. 169–189
- Chrysogonum (Compositae, Heliantheae). Revision of 190-202

- Chrysogonum virginianum 192, var. australe 197, var. virginianum 193
- Chuang, Tsan lang and Lincoln Constance, Cytogeography of Phacelia ranunculacea (Hydrophyllaceae) 115-122
- Coldenia 476; classification of 472
- Coleman, James R. A summary of experimental hybridization in Verbesina (Compositae) 17-31
- Coleochaetaceae in Maine, A new station for 300-304
- Coleochaete pulvinata var. minor var. nov. 302
- Colobogyne: A taxonomic synonym of Spilanthes (Compositae: Heliantheae) 128, 129
- Colt, L. C. Jr., A new station for Coleochaetaceae in Maine 300-304
- Compositae 139-146; genera of the Compositae possessing tetramerous corollas 140-142
- Constance, Lincoln, See Chuang, Tsan lang
- Crompton, C. W., See McNeill, J. and I. J. Bassett
- Cydista aequinoctialis var. hirtella 440 Cytogeography of Phacelia ranunculacea (Hydrophyllaceae) 115-122
- Cytogeography of Sedum lanceolatum and its relatives 95-114
- Disc florets in the Compositae, Observations on tetramerous 139-146
- Distictis buccinatoria 440
- Dorn, Robert D., Willows of the Rocky Mountain states 390-429
- Eastman, L. M., Notable plants of Rattlesnake Mountain, Porter, Maine 316
- Essex County, Massachusetts, Revisions to a flora of 308-315
- Erythronium americanum Ker. (Liliaceae), The pollination ecology of a population of 278-282
- Flora of Gardiners Island 214-239

- Flora of Essex County, Massachusetts, Revisions to a 308-315
- Flora of the Great Dismal Swamp, A contribution towards a vascular 240-268
- Galinsoga (Compositae : Heliantheae), A revision of the genus 319–389
- Galinsoga 319–389; boliviensis sp. nov. 365, caligensis sp. nov. 353, chromosome counts 330 & 331, durangensis comb. nov. 342, elata sp. 340, filiformis 345, formosa nom. nov. 338, glandulosa sp. nov. 380, longipes sp. nov. 348, mandonii 370, mollis 344, parviflora 373, quadriradiata 355, subdiscoidea 368, triradiata sp. nov. 351
- Gardiners Island, Flora of 214-239; checklist of plants 227-239
- Gardner, Robert C., Observations on tetramerous disc florets in the Compositae 139-146
- Gastony, Gerald J., and Douglas E. Soltis, Chromosome studies of Parnassia and Lepuropetalon (Saxifragaceae) from the eastern United States. A new base number for Parnassia 573-578
- Gentry, Alwyn H., Notes on Middle American Bignoniaceae 430-444
- Great Dismal Swamp, A contribution towards a vascular flora of the 240-268
- Grindelia (Compositae-Astereae), A new combination in 583-585
- Grindelia gymnospermoides comb nov. 584
- Grocki, Walter, See Wilce, R. T.
- Heckard, Lawrence and Peter Rubtzoff, Additional notes on Veronica anagallis-aquatica × catenata (Scrophulariaceae) 579-582
- Hehre, Edward J. Jr., The flora of Gardiners Island 214-239
- Hellquist, C. Barre, Observations on some uncommon aquatic plants in

New England 445-451

Hermann, Frederick J. and William A. Weber, Juncus triglumis in North America 160-162

Hybridization in Verbesina (Compositae), A summary of experimental 17– 31

Hybridization between Phyllodoce empetriformis and P. glanduliflora (Ericaceae). An analysis of populations formed by 1–16

Jacaranda copaia ssp. spectabilis comb. et stat. nov. 441

Jansen, Robert and Bruce Parfitt, Allenrolfea mexicana Lundell (Chenopodiaceae): Its conspecificity with A. occidentalis (S. Wats.) Kuntze 130-132

Jones, Samuel B. Jr., See Keeley, Sterling C.

Juncus triglumis in North America

Keeley, Sterling C. and Samuel B. Jones, Jr., Vernonia (Compositae) in the Bahamas – reexamined 147–159

Keil, David J., Chromosome studies in North and Central American species of Pectis L. (Compositae:Tageteae) 70,04

Keil, David J., A revision of Pectis section Pectothrix (Compositae: Tageteae) 32–78

Lauermann, Scott D., See Burk, C. John Levy, Gerald F., See Musselman, L. J. and Daniel L. Nickrent

Lindera benzoin, Red form of 166

Lindera benzoin f. rubra forma nova 166

Lipscomb, Barney L. and Edwin B. Smith, Morphological intergradation of varieties of Bidens aristosa (Compositae) in northern Arkansas 203-213

Lundia puberula 442

McCrea, Kenneth D., See Abrahamson, Warren G.

McNeill, J., I. J. Bassett and C. W. Crompton, Suaeda calceoliformis, the correct name for Suaeda depressa auct. 133–138

Maine, Notable plants of Rattlesnake Mountain, Porter, 316

Maine, A new station for Coleochaetaceae in 300-304

Mansoa verrucifera 443

Massachusetts, Catalpa speciosa naturalized in western 305-307

Massachusetts, Revisions to a flora of Essex County 308-315; [corrigenda 308-311]

Musselman, Lytton J., Daniel L. Nickrent and Gerald F. Levy, A contribution towards a vascular flora of the Great Dismal Swamp 240 268

Najas guadalupensis 450, N. minor 451

New England, Observations on some uncommon aquatic plants in 445-

New England, Rediscovery of Psilocarya nitens in 163, 164

New England, Rudbeckia serotina f. tubuliformis in 165

Nickrent, Daniel L., See Musselman, L., J. and Gerald F. Levy

North America, Battersia mirabilis, A crustose sphacelarioid new to northeastern 292-299

North America, Juncus triglumis in 160-162

North and Central American species of Pectis L. (Compositae : Tageteae). Chromosome studies in 79 94

Notice of publication 167, 317, 465

Pachyptera hymenaea 443, P. standleyi

Parfitt, Bruce, See Jansen, Robert

Parnassia and Lepuropetalon (Saxifragaceae) from the eastern United States. A new base number for Parnassia. Chromosome studies of 573– 578

Pectis L. (Compositae : Tageteae).
Chromosome studies in North and
Central American species of 79-94
Pectis section Pectothrix (Compositae :

Tageteae), A revision of 32-78

Pectis angustifolia 53, var. angustifolia 55, var. fastigiata comb. nov. 60, var. tenella comb. nov. 58; filipes 71, var. filipes 74, var. subnuda 74; incisifolia 63; morphological groups 33; new chromosome counts 82-87, previous chromosome counts 80; papposa 50, var. grandis 53, var. papposa 51; peruviana sp. nov. 48; pringlei 61; purpurea 64, var. purpurea 65, var. lancifolia 68, var. sonorae 67; rusbyi 46; sinaloensis 69; stenophylla 39, var. biaristata comb. nov. 45, var. gentryi var. nov. 42, var. puberula comb. nov. 44, var. rosei comb. nov. 44, var. stenophylla 42; uniaristata var. jangadensis comb. nov. 90; vollmeri 75

Phacelia ranunculacea (Hydrophyllaceae), Cytogeography of 115-122

Phyllodoce empetriformis and P. glanduliflora (Ericaceae), An analysis of populations formed by hybridization between 1-16

Pithecoctenium crucigerum 443

Pollination ecology of a population of Erythronium americanum Ker. (Liliaceae) 278-282

Poole, James P., Revisions to a flora of Essex County, Massachusetts 308-315

Porter, Duncan M., Typification of Tiquilia darwinii and Tiquilia fusca (Boraginaceae) 288–291

Potamogeton filiformis 445, hillii 446, lateralis 450, × longiligulatis 448, yaginatus 446

Pseudocatalpa caudiculata 443

Psilocarya nitens in New England, Rediscovery of 163, 164 Psilostrophe DC. (Compositae) I. Chromosome variability. Biosystematics of 169–189

Psilostrophe 169–189; chromosome configurations in experimental hybrids of 182; chromosome variability 187; meiotic configurations in natural populations of 174–177

Quercus × introgressa, A new hybrid Oak 453-464

Quercus × introgressa hybrida nova 453

Richardson, Alfred T., Monograph of the genus Tiquilia (Coldenia, sensu lato), Boraginaceae : Ehretioideae 467-572

Rocky Mountain States, Willows of the 390-429

Rubtzoff, Peter, See Heckard, Lawrence Rudbeckia (Compositae), Ultraviolet light reflection and absorption patterns in populations of 269-277

Rudbeckia serotina f. tubuliformis in New England 165

Ruffin, John, A new combination in Grindelia (Compositae-Astereae) 583-585

Saccharum or Saccarophorum? 123-127

Salix 390-429; alba 405; amygdaloides 403; arctica 412; arizonica 424; babylonica 406; barclayi 420; barrattiana 427; bebbiana 413; bonplandiana 402; boothii 423; brachycarpa 411; candida 418; cascadensis 413; commutata 421; discolor 415; drummondiana 417; eastwoodiae 422; exigua 407; farriae 420; fragilis 406; geveriana 414; glauca 411; gooddingii 402; irrorata 418; lasiandra 405, var. caudata 405; lasiolepis 419; lemmonii 415; ligulifolia 426; lucida 404; lutea 425; melanopsis 408; monochroma 426; monticola 424; orestera 422; pedicellaris 410; pentandra 403; petiolaris 414; planifolia 416; pseudomonticola 419; reticulata ssp. nivalis 409; rotundifolia 410; scouleriana 416; serissima 404; sitchensis 427; taxifolia 406; tweedyi 428; vestita 408; wolfii 421

Sedum lanceolatum and its relatives. Cytogeography of 95-114

Shaw, Elizabeth A., Saccharum or Saccarophorum? 123-127

Smith, Edwin B., See Lipscomb, Barney L.

Soltis, Douglas E., See Gastony, Gerald J.

Sorrie, Bruce A., Rediscovery of Psilocarya nitens in New England 163, 164

Spilanthes (Compositae: Heliantheae), Colobogyne: A taxonomic synonym of 128, 129

Spilanthes langbianensis comb. nov.

Stigmatic surfaces in the Begoniaceae, Preliminary notes on the 283–287 Stizophyllum riparium 443

Stuessy, Tod F., Colobogyne: A taxonomic synonym of Spilanthes (Compositae: Heliantheae) 128, 129

Stuessy, Tod F., Revision of Chrysogonum (Compositae : Heliantheae) 190-202

Suaeda calceoliformis, The correct name for Suaeda depressa auct. 133-138

Tabebuia donnell-smithii 443, T. impetiginosa 443

Tetramerous disc florets in the Compositae, Observations on 139-146

Thomson, Paul M., Quercus × introgressa, A new hybrid Oak 453-464

Tiquilia atacamensis 551; canescens 506, var. canescens 508, var. pulchella 509; classification of 474; conspicua 559; chromosome number, species examined for 481-486; cuspidata 537; darwinii 566; dicho-

toma 542; elongata 554; ferreyrae 549; **galapagoa** comb. nov. 563; gossypina 525; grandiflora 547; greggii 510; hispidissima 517; hunteri 556; latior 515; litoralis 557; mexicana 524; nesiotica 565; nuttallii 532; palmeri 538; paronychioides 561; plicata 534; purpusii 529; simulans 545; tacnensis 553; tuberculata 519; turneri 521

Tiquilia, (Coldenia, sensu lato), Boraginaceae: Ehretioideae, Monograph of the genus 467-572

Figuilia darwinii and Tiquilia fusca (Boraginaceae), Typification of 288– 291

Uhl, Charles H., Cytogeography of Sedum lanceolatum and its relatives 95-114

Ultraviolet light reflection and absorption patterns in populations of Rudbeckia (Compositae) 269-277

United States. A new base number for Parnassia. Chromosome studies of Parnassia and Lepuropetalon (Saxifragaceae) from the eastern 573-578

Verbesina (Compositae), A summary of experimental hybridization in 17-31

Verbesina alternifolia × helianthoides 23: aristata × glabrata 18: heterophylla × occidentalis 18: lindheimeri × occidentalis 20: lindheimeri × rothrockii 19: virginica × heterophylla 19: walteri × alternifolia 24: walteri × occidentalis 23

Vernonia (Compositae) in the Bahamas reexamined 147-159

Vernonia arbuscula 156; blodgettii 157; characteristics to distinguish some species 148; cinerea 158; fertility in species and hybrids 155; leaf shapes 152

Veronica anagallis-aquatica × catenata (Scrophulariaceae), Additional notes on 579-582

- Watson, Thomas J. Jr., An analysis of populations formed by hybridization between Phyllodoce empetriformis and P. glanduliflora (Ericaceae) 1-
- Weber, William A., See Hermann, Frederick J.
- Wilce, R. T., and Walter Grocki, A crustose sphacelarioid new to northeastern North America: Battersia mirabilis 292-299
- Willows of the Rocky Mountain States 390-429

